

# Nutritional values and Chemical Constituents of *Anadara brughtonii* (Schrenck, 1867) and *Crassostea gigas* (Thunberg, 1793) used in Traditional Medicine in Myanmar

Thein Gi Naing<sup>1</sup>, Hnin Hnin Htun<sup>1</sup>, Win Soe<sup>1</sup>, Than Tun<sup>1</sup>,  
Maung Maung Thet<sup>1</sup>, Khin Mya Mya<sup>2</sup>, Thant Zin<sup>2</sup>,  
Thein Zaw Lin<sup>1</sup>

1 University of Traditional Medicine, Mandalay ,  
2 Department of Zoology, Mandalay University

# INTRODUCTION

- ❖ Myanmar used animals and their products as medicinal substances since ancient times.
- ❖ Most of Myanmar traditional drugs are derived from sources of plants and animals.
- ❖ Wild and domestic animals and their by-products (e.g., hooves, skins, bones, feathers, milk and tusks) form important ingredients in the preparation of curative, protective and preventive medicine.

- ❖ According to the World Health Organization (1993), about 80% of the world people rely primarily on animal and plant-based medicines.
- ❖ About 20 % of Myanmar traditional medicine is based on animal-derived substances.

# LITERATURE REVIEW

# *Anadara broughtonii*

- ❖ species of Ark clam.
- ❖ In the Myanmar costal water it had been recorded from Ngapali, Maungmagan, KyaukKalat., Cocos Island.
- ❖ The species is distributed in Far East, from Russia down to Korea, Mainland China, Japan and Taiwan(Cho et.al.,2007)



# Traditional medicine in Myanmar

- shell of *Anadara broughtonii* was also used by rulers of early Myanmar dynasties.
- The ash form of Jau thwa khone has given to remove toxin from the body,
- increased energy power, salty and cool in nature.

- ❖ It also used in the treatment of
- ❖ indigestion,
- ❖ certain gastric and intestinal disorder,
- ❖ cough,
- ❖ asthma,
- ❖ chest injuries,
- ❖ mouth disease,
- ❖ oliguria,

- ❖ dysentery,
- ❖ hotness of urine,
- ❖ piles,
- ❖ bowel disorder,
- ❖ blood vomiting and menstrual disorders in Myanmar traditional medicine (AshinNagathein, 1972).



# *Crassostrea gigas*

- ❖ The Pacific oyster *Crassostrea gigas* is a marine invertebrate
- ❖ belonging to the family Ostreidae (Mollusca, Bivalvia)
- ❖ worldwide distribution from Japan to occidental countries in Europe and America (Guoet al., 2008)



# Traditional medicine in Myanmar

- ❖ Shell of *Crassostrea gigas* was used by rulers of early Myanmar dynasties.
- ❖ Traditional medicine, As the In terms of Myanmar Traditional Medicine, taste of ka nu ka mar is sweet and cool in nature.
- ❖ Recommended dosage is 1 to 2 grams of ash.
- ❖ The ash of ka nu ka mar has been given promote digestive power and stimulate appetite

- ❖ It is also used in the treatment of
- ❖ indigestion,
- ❖ hepatitis,
- ❖ certain gastric and intestinal disorder,
- ❖ cough, asthma,
- ❖ chest injuries,
- ❖ mouth disease,
- ❖ oliguria,
- ❖ dysentery, bowel disorder, blood vomiting and menstrual disorders in Myanmar traditional medicine (AshinNagathein, 1972).

# Traditional Medicine Formulae

- ❖ which are commonly used in Myanmar Traditional Medicine formulation (TMF).
- ❖ Jau thwa khone pyar (Ash) is the ingredient of TMF6 (Ha Leik da Sonna Ngan Hsei).
- ❖ Ka nu ka mar is the ingredients of TMF-24 Na Ga Ra (Lei;Hnjin;KaLa'Hsei;).

- ❖ Though these samples are commonly in Myanmar Traditional Medicine much work has not been reported on the proximate and chemical compositions.
- ❖ Therefore these crude animal's products were selected for chemical and pharmacological investigation

# OBJECTIVES

- ❖ to identify the two species of shells that are used in Myanmar Traditional Medicine
- ❖ to investigate the nutritional values and chemical constituents of two species of shells used in Myanmar Traditional Medicine

# **METHODOLOGY**

# Study Design

❖ Zoological identification of crude drugs.



# Study Area



# Study Period

- from July 2018 to February 2019.

# Materials and Methods

- ❖ Extraction of samples is one of the procedure of Association of Official Analytical Chemistry (A.O. A. C).
- ❖ Determination of elements by Energy Dispersive X-ray Fluorescence Spectrophotometer (EDXRF).

# Statistical Analysis

- The results were expressed as mean  $\pm$  standard deviation (SD).

# Samples Collection

- ❖ Shells of *A. broughtonii* and *C. gigas* were purchased locally from Baja Hsei: zain(traditional medicine shop)
- ❖ it had been recorded from Ngapli, Maungmagan, KyaukKalat, Coco Island locations in Myanmar Coastal water
- ❖ Specimen were identified according to the Abbott, R.T. (1991).



# Samples Preparation




# Samples Preparation

- ❖ The samples 150g shells of *Anadara broughtonii* and *Crassostrea gigas* were first washed thoroughly with distilled water
- ❖ washed with sterile water mixed with 3cc lime juice
- ❖ washed with sterile water to remove foreign matters
- ❖ sample is dried at drying condition in oven.


- ❖ One hundred and fifty grams of samples were crushed into smaller pieces and then make powder by blender.
- ❖ The powder was sieved using a stainless steel sieve to get fine powder and
- ❖ then sterilized for an hour in the oven at 105°C and stored in bottles prior to analysis.
- ❖ Proximate analysis, mineral composition and pharmaceutical product were carried out on dried powder.




# Preparation of Powder and Ash form




A. Powder of *Anadara broughtonii*



B. Ash of *Anadara broughtonii*



C. Powder of *Crassostrea gigas*



D. Ash of *Crassostrea gigas*

# RESULTS

# Systematic Position of the Studied Species

Phylum	-	Mollusca
Class	-	Bivalvia
Order	-	Arcida
Family	-	Arcidae
Genus	-	<i>Anadara</i>
Species	-	<i>A.broughtonii</i>
Order	-	Ostreoida
Family	-	Ostreidae



Genus - *Crassostrea*

Species - *C. gigas*

*Crassostrea gigas*

Synonym

*Magallana gigas* (Thunberg, 1793)

*Anadara broughtonii* (Schrenck, 1867)



# Description of Study Species

## 1. Shell of *Anadara broughtonii*

- Scientific name - *Anadara broughtonii*
- Local name - Jau thwa khone
- Common name - Sea mussel





A. Dorsal view of shell of  
*Anadara broughtonii*.



B. Ventral view of shell of  
*Anadara broughtonii*.

- ❖ Shell of *Anadara broughtonii* - 6 to 9 cm length
- ❖ It is thick, shell equivalve, solid, ovate, strongly inflated, slightly longer than high and feebly in equilateral.
- ❖ About 18 radial ribs (15 to 20) with wide interstices at each valve.
- ❖ Ribs stout and distinctly rugose, bearing regular, often rectangular nodules.



- ❖ Internal margins with strong crenulations corresponding with the external radial ribs.
- ❖ Outside of shell white under the yellowish brown periostracum.
- ❖ Inner side white, often tinged light yellow towards the umbonal cavity (Plate 3).





## 2.Shell of *Crassostrea gigas*

- ❖ Scientific name - *Crassostrea gigas*
- ❖ Local name - ka nu ka mar
- ❖ Common name - Oyster
- ❖ The shell of *Crassostrea gigas* is large, rounded, radial folds are often extremely rough and sharp.



- ❖ The two valves of the shell are slightly different in size and shape, the right valve being moderately concave.
- ❖ Shell colour is variable, usually pale white or off-white. Mature specimens can vary from 80 to 400 mm long (Plate4).




. Ventral view of right valve  
of *Crassostrea gigas*




Ventral view of left valve of  
*Crassostrea gigas*




A. Dorsal view of right valve of  
*Crassostrea gigas*



B. Ventral view of right valve of  
*Crassostrea gigas*



C. Dorsal view of left valve of  
*Crassostrea gigas*



D. Ventral view of left valve of  
*Crassostrea gigas*

# Table 1 Proximate composition of powder of shells of *Anadara broughtonii*

No	Parameters	Value 1 (%)	Value 2 (%)	Value 3 (%)	Value 4 (%)	Value 5 (%)	Mean $\pm$ SD (%)
1	Moisture	0.28	0.27	0.26	0.28	0.28	0.28 $\pm$ 0.01
2	Ash	97.65	97.75	97.65	98.45	98.45	97.75 $\pm$ 0.04
3	Carbohydrate	0.00	0.00	0.00	0.00	0.00	0 $\pm$ 0.00
4	Protein	0.16	0.17	0.19	0.5	0.5	0.2 $\pm$ 0.01
5	Fiber	2.48	2.46	2.47	2.47	2.48	2.47 $\pm$ 0.01
6	Fat	0.73	0.72	0.73	0.72	0.73	0.73 $\pm$ 0.01

**Table 2 Total ash, water soluble ash and acid insoluble ash content of powder of shells of *Anadara broughtonii* .**

---

<b>No.</b>	<b>Parameters</b>	<b>Quantity (%)</b>
<hr/>		
1	Total ash	97.75±0
2.	Water soluble	3.7 ±0
3.	Acid insoluble	76.75±0

---

**Table 3 Percentage of macroelements involved  
in powder of shells of *Anadara broughtonii***

No	Elements	Quantity 1 (%)	Quantity 2 (%)	Quantity 3 (%)	Mean±SD (%)
1	Magnesium (Mg)	1.09	1.09	1.09	1.1±0.1
2	Sodium (Na)	0.00	0.00	0.00	0.00±0
3	Calcium (Ca)	52.75	52.75.	52.75	52.75±0
4	Chlorine (Cl)	0.00	0.00	0.00	0.00±0
5	Potassium (K)	0.00	0.00	0.00	0.00±0
6	Sulfur (S)	0.00	0.00	0.00	0.00±0

# Table 4 Percentage of microelements involved in Powder of shells of *Anadara broughtonii*

No	Elements	Quantity 1 (%)	Quantity 2 (%)	Quantity 3 (%)	Mean±SD (%)
1	Aluminium (Al)	-	-	-	-
2	Silicon (Si)	0.2	0.2	0.2	0.2±0
3	Manganese (Mn)	0.1	0.1	0.1	0.1±0.01
4	Iron (Fe)	0.01	0.01	0.01	0.1±0
5	Copper (Cu)	0.0002	0.0002	0.0002	0.0002±0
6	Zinc (Zn)	0.01	0.01	0.01	0.01±0

**Table 5 Percentage of heavy metals involved in powder of shells of *Anadara broughtonii***

No	Elements	Quantity 1 (%)	Quantity 2 (%)	Quantity 3 (%)	Mean±SD (%)
1	Arsenic (As)	-	-	-	-
2	Cadmium (Cd)	0.0003	0.0003	0.0003	0.0003±0
3	Mercury (Hg)	-	-	-	-
4	Lead (Pb)	0.0008	0.0008	0.0008	0.0008±0



# Table 6 Proximate composition of powder of shells of *Crassostrea gigas*

No	Parameters	Value 1 (%)	Value 2 (%)	Value 3 (%)	Value 4 (%)	Value 5 (%)	Mean $\pm$ SD (%)
1	Moisture	0.27	0.25	0.24	0.24	0.24	0.24 $\pm$ 0.01
2	Ash	97.15	97.16	97.25	97.15	99.36	97.16 $\pm$ 0.1
3	Carbohydrate	0	0	0	0	0.17	0 $\pm$ 0.00
4	Protein	0.58	0.57	0.56	1	1	0.58 $\pm$ 0.24
5	Fiber	1.92	1.93	1.95	1.93	1.93	1.93 $\pm$ 0.01
6	Fat	0.54	0.56	0.55	0.09	0.09	0.54 $\pm$ 0.3

**Table 7 Total ash, Water soluble ash and Acid insoluble ash Contents of powder of shells of *Crassostrea gigas***

<b>No.</b>	<b>Parameters</b>	<b>Quantity((%)</b>
1.	Total ash	97.16±0.014
2.	Water soluble	1.75±0
3.	Acid insoluble	72.44±0

**Table 8 Percentage of macroelements involved  
in Powder shells of *Crassostrea gigas***


No	Elements	Value 1 (%)	Value 2 (%)	Value 3 (%)	Mean $\pm$ SD (%)
1.	Magnesium(g)	3.2311	3.2311	3.2311	3,2311 $\pm$ 0
2.	Sodium (Na)	0.00	0.00	0.00	0.00 $\pm$ 0
3.	Calcium (Ca)	54	54	54	54 $\pm$ 0
4.	Chlorine (Cl)	0.00	0.00	0.00	0.00 $\pm$ 0
5.	Potassium (K)	0.00	0.00	0.00	0.00 $\pm$ 0
6.	Sulfur (S)	0.00	0.00	0.00	0.00 $\pm$ 0

**Table 9 Percentage of microelements involved  
in Powder of shells of *Crassostrea gigas***


No	Elements	Value 1 (%)	Value 2 (%)	Value 3 (%)	Mean $\pm$ SD (%)
1	Aluminium (Al)	0.00	0.00	0.00	0.00 $\pm$ 0
2	Silicon (Si)	0.1	0.1	0.1	0.1 $\pm$ 0
3	Manganese(Mn)	0.2	0.2	0.2	0.2 $\pm$ 0
4	Iron (Fe)	0.13	0.13	0.13	0.13 $\pm$ 0
5	Copper (Cu)	0.0004	0.0004	0.0004	0.0004 $\pm$ 0
6	Zinc (Zn)	0.2	0.2	0.2	0.2 $\pm$ 0

**Table10 Percentage of heavy metals involved  
in Powder of shells of *Crassostrea gigas***


No	Elements	Quantity	Quantity	Quantity	Mean±S
		1 (%)	2 (%)	3 (%)	D
1	Arsenic (As)	0.0001	0.0001	0.0001	0.0001
2	Cadmium(Cd)	-	-	-	-
3	Mercury (Hg)	-	-	-	-
4	Lead (Pb)	0.0005	0.0005	0.0005	0.0005



**Fig. 1 Proximate Composition of Powder of shells of *Anadara broughtonii***




**Fig.2 Percentage of macroelements involved in powder of shells of *Anadara broughtonii***




**Fig. 3 Percentage of microelements involved in Powder of shells of *Anadara broughtonii***







**Fig.4 Total ash, water soluble ash and acid insoluble ash content of powder of shells of *Anadara broughtonii***




**Fig. 5 Percentage of heavy metals involved in powder of shells of *Anadara broughtonii***




**Fig. 6 Proximate composition of powder of shells of *Crassostrea gigas***




**Fig.7 Percentage of macroelements involved in powder of shells of *Crassostrea gigas***



**Fig. 8 Percentage of microelements involved in powder shells of *Crassostrea gigas***



**Fig. 9 Total ash, water soluble ash and acid insoluble ash contents of *Crassostrea gigas***



**Fig. 10 Percentage of heavy metals involved in powder of shells of *Crassostrea gigas***

# DISCUSSION

- The Proximate Composition of shells Powder of *Anadara broughtonii* and *Crassostrea gigas* revealed that these contained reasonable amounts of moisture, carbohydrate, fiber, protein and fat.
- Carbohydrate and fat contents were found to be very low in these samples. This confirms that sample is not a good source of fat.



- The contents of fiber were found to be considerable amount in these samples.
- Mixtures of soluble and insoluble fibers improve diabet-ic glucose control and lower serum triglycerides (Anderson 1990).
- Protein provides essential amino acids, particularly important during growth and development, and it is a source of energy (Thomas *et al.*, 2004).

- The content of ash was found to be highest in these samples (Table 1,6 and Fig 1,6).
- It is a reflection of total inorganic matter present in these samples and also indicates that they possess the most abundant mineral like calcium, which are essential for good health (Oloyede, 2008).

- In this study cadmium, lead and mercury were found to contain below levels according to the WHO guideline values of cadmium, lead and mercury.
- Arsenic and mercury were not found in powder of *Anadara broughtonii*.
- Cadmium and mercury were not found in powder *Crassostrea gigas* .
- This finding indicated that these animal parts may be used safely in the traditional medicine.

# ACKNOWLEDGEMENT

- Dr Moe Swe, Director General, Department of Traditional Medicine, Naypyidaw.
- Dr Thant Zin Professor and Head, Department of Zoology, University of Mandalay
- Dr Thein Zaw Lin, Actinng Rector, University of Traditional Medicine, Mandalay for their kind to conduct this present study.

- Dr Khin Mya Mya, Professor, Department of Zoology, University of Mandalay, her patient guidance, kind supervision and constant encouragement in my research work.
- I am gratefully indebted to Dr Hla Swe, Head, Department of Academic Affairs, Retired at University of Yangon for his invaluable advice in my research work.
- I am also thank to Dr. Moe Kyaw Myint, Deputy Director/Head, Parasitology Research Division, Department of Medicinal Research, Pyin OoLwin Branch.
- I am gratefully indebted to my parents U Ye Naing and Daw Khin Saw Yi for their kindness and encouragement in my research work.

# REFERENCES

- A.O.A.C., 2000. *Official methods of Analysis* 17th Edn. Washington DC: Association of official Analytical Chemist, Washington. pp 920-978
- Adverse Health Effects Of Heavy Metals In Children, 2011. Children's Health and the Environment WHO Training Package for the Health Sector World Health Organization. Available from: [www.who.int/ceh](http://www.who.int/ceh) (Accessed August 2016).
- American Herbal Products Association, 2009. Heavy metal analysis and limits in herbal dietary supplement
- AshinNagathein, 1972. Jauthwakhone, *JA Tha Ta Jinge Ni DaThamoudajaKjan:Gji*, Han Thawati Beidaga Poun Hnei Tai, Yangon, Myanmar. 211-213.
- AshinNagathein, 1972. ka nu ka mar, *JA Tha Ta Jinge Ni DaThamoudajaKjan:Gji*, Han Thawati Beidaga Poun Hnei Tai, Yangon, Myanmar. 210-211.

***THANK YOU***