Minerals Evaluation, Antibacterial Activities and Antioxidant Activities of Zingiber officinale Roscoe (ချင်း)

Lei LeiWin¹,KyawtKyawtKhaing¹,Khin Lay Sein¹, Nu Ye Thin¹,Aye Thi¹, Saw MyatThwe¹, Cheery Kyawwin¹,NweNweYi²,KhaingKhaingMar¹ and Win Aung¹

1. Department of Medical Research (PyinOoLwin Branch)

2. Department of Biology (Sagaing University of Education)

The medicinal value of plants have assumed important dimension in the past few decades owing largely to the discovery that extracts from plants contain not only minerals and primary metabolites but also adiverse array of secondary metabolites with antioxidant and antimicrobial potentials.Owing to its minimal toxicity, cost effectiveness and pharmacologically active, 80% of population in developing countries depend on traditional medicines, mostly natural plant products, for their primary health care needs as estimated by World Health Organization. A number of minerals essential to human nutrition are accumulated in different parts of plants. The alarming increase in the rate of infection by antibiotic resistant microorganisms has urged scientists to search for compounds which have potential antimicrobial activity. Antioxidants are vital substances which possess ability to protect the body from damage caused by free radical induced oxidative stress. The purpose of current study was mineral evaluation, antimicrobial activities and antioxidant activities of ZingiberofficinaleRoscoe, ginger(csif;).Atomic absorptionspectrophotometerand UV-Visible spectro- photometer were used for determination of minerals and antioxidant activities. Antibacterial activity was determined by agar disc diffusion method(WHO, 2003).Broth dilution method was used for determination of Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC). Macrominerals content of ginger were within permissible limit of Ajasa, 2004 and content of microminerls were within permissible limit of vegetables set byFAO/WHO, 2001.Ginger extracts showed zone of inhibition12 mm – 15mm against Pseudomonas aeruginosa,7mm – 9 mm against Staphylococcus aureus and 12 mm – 15mm against Escherichia coli. Although MIC and MBC of aqueous extracts did not show the concentration up to 10mg/mL, MIC of ethanolic extract was observed in the range from 1mg/mL to 8mg/mL and MBC from 2mg/mL to 9mg/mL. The antioxidant activity of aqueous extract of ginger (IC₅₀: 0.022ug/ml) and ethanolic extract (IC₅₀: 0.333ug/ml)were comparable to well-known antioxidant, ascorbic acid (IC₅₀: 0.59) ug/ml). Thus, ginger has strong free radical scavenging activity. These findings indicated that, ginger has potential to provide nutrients to human beings, preventive properties against oxidative damage and some pathogens. These results willgive scientific information for herbal medicine users, local practitioners, standardization &quality control of precious indigenous drug and pharmaceutical industries using ginger for different types of ailments.

Key Words: Macrominerals, Microminerals, Antibacterial activity, Antioxidant activity, *Zingiber officinale* Roscoe.